

Short report

Open Access

Testosterone correlates with Venezuelan equine encephalitis virus infection in macaques

Michael P Muehlenbein^{*1}, Frank B Cogswell², Mark A James³, James Koterski⁴ and George V Ludwig⁴

Address: ¹Department of Anthropology, University of Wisconsin-Milwaukee, USA, ²Department of Bacteriology and Parasitology, Tulane National Primate Research Center, Covington, Louisiana, USA, ³Department of Tropical Medicine, Tulane University School of Public Health and Tropical Medicine, New Orleans, Louisiana, USA and ⁴Diagnostic Systems Division, United States Army Medical Research Institute of Infectious Diseases, Ft. Detrick, Maryland, USA

Email: Michael P Muehlenbein^{*} - mpm1@uwm.edu; Frank B Cogswell - cogswell@tpc.tulane.edu; Mark A James - mjames@tulane.edu; James Koterski - James.Koterski@amedd.army.mil; George V Ludwig - George.Ludwig@amedd.army.mil

^{*} Corresponding author

Published: 29 March 2006

Received: 06 September 2005

Virology Journal 2006, **3**:19 doi:10.1186/1743-422X-3-19

Accepted: 29 March 2006

This article is available from: <http://www.virologyj.com/content/3/1/19>

© 2006 Muehlenbein et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Here we briefly report testosterone and cytokine responses to Venezuelan equine encephalitis virus (VEEV) in macaques which were used as part of a larger study conducted by the Department of Defense to better characterize pathological responses to aerosolized VEEV in non-human primates. Serial samples were collected and analyzed for testosterone and cytokines prior to and during infection in 8 captive male macaques. Infected animals exhibited a febrile response with few significant changes in cytokine levels. Baseline testosterone levels were positively associated with viremia following exposure and were significantly higher than levels obtained during infection. Such findings suggest that disease-induced androgen suppression is a reasonable area for future study. Decreased androgen levels during physiological perturbations may function, in part, to prevent immunosuppression by high testosterone levels and to prevent the use of energetic resources for metabolically-expensive anabolic functions.

Findings

Venezuelan equine encephalitis virus (VEEV) complex, like the Western and Eastern equine encephalitis virus complexes, is a collection of alphaviruses (single, positive strand RNA) of the family *Togaviridae* that infect humans, rodents and equines in the Americas, and are transmitted via various arthropods, including mosquitoes such as *Aedes*, *Culex*, and *Psorophora* [1]. VEEV is a neurotropic virus in laboratory rodents that induces illness characterized first by replication in the lymphoid tissues and development of a high viremia, and second by invasion of the central nervous system via the olfactory neuroepithelium resulting in encephalitis [2-4]. The murine immune

response to VEEV is characterized by both Th-1 and Th-2 cytokine gene expression, and both antibody- and cell-mediated responses may play important roles in protection against VEEV [5,6]. Among human and non-human primates, the virus is much less fatal, usually resulting in fever with viremia and lymphopenia [7]. The specific immune responses to VEEV in humans or non-human primates are not well known.

Here we report preliminary results describing the testosterone and cytokine responses to VEEV in macaques which were used as part of a larger study conducted by the Department of Defense to better characterize pathological

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 29 MAR 2006		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Testosterone correlates with Venezuelan equine encephalitis virus in macaques, Virology Journal 3:19 - 23				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Muehlenbein, MP Cogswell, FB James, MA Koterski, J Ludwig, GV				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) United States Army Medical Research Institute of Infectious Diseases, Fort Detrick, MD				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Here we briefly report testosterone and cytokine responses to Venezuelan equine encephalitis virus (VEEV) in macaques which were used as part of a larger study conducted by the Department of Defense to better characterize pathological responses to aerosolized VEEV in non-human primates. Serial samples were collected and analyzed for testosterone and cytokines prior to and during infection in 8 captive male macaques. Infected animals exhibited a febrile response with few significant changes in cytokine levels. Baseline testosterone levels were positively associated with viremia following exposure and were significantly higher than levels obtained during infection. Such findings suggest that disease-induced androgen suppression is a reasonable area for future study. Decreased androgen levels during physiological perturbations may function, in part, to prevent immunosuppression by high testosterone levels and to prevent the use of energetic resources for metabolically-expensive anabolic functions.					
15. SUBJECT TERMS Venezuelan equine encephalitis virus, VEE, testosterone response, cytokine response, laboratory animals, nonhuman primates					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

responses to aerosolized VEEV in non-human primates. Ample evidence suggests that testosterone is inversely associated with measures of immunity in several species, including primates [8-11]. It was therefore hypothesized that testosterone would be directly associated with VEEV viremia and inversely associated with proinflammatory cytokine levels.

Eleven captive-born, adult (mean age: 9.74 years) male cynomolgus monkeys (*Macaco fascicularis*) were maintained in individual steel cages in a BSL-3 facility at the United States Army Medical Research Institute of Infectious Diseases (USAMRIID), Fort Detrick, Maryland. Animals were maintained on a standard laboratory primate diet, and water was available ad libitum. All animals tested negative for antibodies to Western, Eastern, and Venezuelan Equine Encephalitis viruses before their use in this experiment.

Blood samples were obtained on pre-exposure days -60, -45, and -30. On pre-exposure day -30, each animal was implanted with a Data Sciences International TA11PA-C40 Temperature and Activity Telemetry Transmitter. On day 0 (day of exposure), 8 randomly chosen animals were exposed to a total inhaled dose of 1×10^8 plaque forming units (PFU) of VEEV, Trinidad Donkey strain. Every 24 hours post-exposure, 2 of the 8 infected animals were randomly chosen to be anesthetized and euthanized by exsanguination. Euthanasia and single interval sampling, rather than multiple bleeds throughout infection, were the requirement of another study by the Department of Defense using these same animals.

For each blood sample, animals were anesthetized with 0.1 mg/kg ketamine/acepromazine maleate solution and a 6 ml blood sample was immediately obtained from the femoral vein using a Vacutainer tube with serum separator. Samples were centrifuged, the serum frozen at -80°C , and aliquots shipped to Yale University and Tulane University. Samples from the two animals which were exposed to VEEV and sampled/sacrificed 24 hours later (day 1) were not available for analysis in the current study, thus the final size was as follows: 3 baseline samples taken from each of 11 animals ($N = 33$); 1 sample taken from each of 2 animals 48 hours after exposure ($N = 2$); 1 sample taken from each of 2 animals 72 hours after exposure ($N = 2$); 1 sample taken from each of 2 animals 96 hours after exposure ($N = 2$). Because of ethical concerns and costs associated with the use of non-human primates, and because this project utilized animals from a study in progress by the Department of Defense, sample sizes could not be increased for purposes described here.

Research was conducted in compliance with the Animal Welfare Act and other Federal statutes and regulations

relating to animals and experiments involving animals and adheres to principles stated in the Guide for the Care and Use of Laboratory Animals, National Research Council, 1996. The facility where this research was conducted is fully accredited by the Association for Assessment and Accreditation of Laboratory Animal Care International.

At Yale University, samples were assayed for total testosterone and cortisol levels using coated-tube radioimmunoassay kits (DSL-4000 and DSL-2100) from Diagnostic Systems Laboratories (DSL), Webster, Texas. The sensitivities of the assays were 0.08 ng/ml for testosterone and 0.3 ng/dl for cortisol. The correlation coefficients for each of the curves were better than 0.99. High and low level DSL controls were included in each standard curve; results for the controls in each assay were within established confidence limits. Intra-assay coefficients of variation were assessed using the mean coefficients of variation of control duplicates. Intra-assay coefficients of variation were 5.8% for testosterone and 9.1% for cortisol. All samples were run in a single assay, therefore there were no inter-assay coefficients of variation to assess. For each individual animal, hormone values from the pre-exposure samples on days -60, -45, and -30 were averaged to yield a baseline (pre-exposure) level.

At Tulane University, samples were assayed for the following cytokines using enzyme-linked immunosorbent assay: IL-1 β , IL-4, IL-10, IL-12, TNF α , and IFN γ (Biosource International Inc., Camarillo, CA, USA). Of each of these kits, all but IL-1 β (human IL-1 β kit) was designed specifically for use in macaques. The human IL-1 β kit demonstrates 100% cross-reactivity with non-human primate samples (S. Durham, BioSource International, Technical Services, personal communication). The sensitivities of the assays were: IFN γ : <4 pg/ml; TNF α : <2 pg/ml; IL-1 β : 1 pg/ml; IL-4: <3 pg/ml; IL-10: <10 pg/ml; IL-12: <4 pg/ml. The correlation coefficients for each of the curves were better than 0.96. Intra-assay coefficients of variation were assessed using the mean coefficients of variation of control duplicates. Intra-assay coefficients of variation were: IFN γ : 5.1%; TNF α : 3.7%; IL-1 β : 4.7%; IL-4: 5.4%; IL-10: 5.1%; IL-12: 4.3%. All samples were run on a single assay, and therefore there were no inter-assay coefficients of variation to assess.

At USAMRIID, viremia was determined by titration of samples on tissue culture (VERO 76) cells and enumeration of plaques (plaque forming units: PFUs). Viremia was measured only on the day of sacrifice, and therefore this measure does not reflect peak viremia titers.

Data were analyzed using SAS/STAT software (SAS Institute Inc., Cary, NC). The Wilcoxon Signed Rank test was used to test whether the median change in temperature,

Table 1: Viremia and pre- and post-infection testosterone levels for each of six animals infected with VEEV

Animal ID	Viremia (PFU) at time of sacrifice	Time (hrs) between exposure and sacrifice	Testosterone (ng/ml) at time of sacrifice	Pre-exposure testosterone (ng/ml) (mean for each animal)
1	7800	48	15.58	28.14
2	7800	48	20.43	26.29
3	370	72	22.06	16.44
4	370	72	2.81	13.11
5	50	96	12.56	25.07
6	25	96	6.79	8.01

testosterone, cortisol, or cytokine values from the pre-exposure samples to the sacrifice (exposure) samples were significantly different from zero (no change). Spearman correlations were used to determine associations between the various variables. Cortisol levels were determined and controlled for in (as a covariate) all analyses involving testosterone because cortisol exhibits mainly inhibitory effects on immune function and because cortisol is often associated with suppression of the hypothalamic-pituitary-testicular axis [12,13]. As the immunosuppressive actions of glucocorticoids are better characterized than that of androgens, the focus of this study was testosterone rather than cortisol-related effects. Additionally, level of significance (α) was set at 0.10 so as to err on the side of conservatism (to avoid false negatives) given the small sample size. Power analyses for the Wilcoxon Signed Rank tests were performed with the SAS/STAT Analyst Sample Size application for paired t-tests (SAS Institute Inc., Cary, NC) to assess the likelihood of false negatives. Power (%) is reported below for those tests that revealed non-significant results.

Temperature, virus titer, testosterone and cytokine levels were compared between samples taken prior to exposure and on the day of sacrifice (48, 72, or 96 hours post-exposure, depending on animal). Mean temperature at time of sacrifice (37.25°C) was significantly higher ($T = 10.5$, $p = 0.03$) than mean temperature prior to VEEV exposure (35.89°C). Mean testosterone level at time of sacrifice (13.37 ng/ml) was significantly lower ($T = -8.5$, $p = 0.094$) than testosterone level prior to VEEV exposure (19.51 ng/ml). Table 1 presents viremia and pre-and post-infection testosterone levels for each of the six animals infected with VEEV.

Statistical analyses could only be performed on IL-12 and IL-10 levels because TNF α and IL-4 levels were not above a detectable level for any sample of any animal, and only one animal had IL-1 β and IFN γ values greater than zero. Mean IL-12 prior to exposure was 537.36 pg/ml , which was significantly higher ($T = -10.5$, $p = 0.03$) than mean IL-12 at time of sacrifice (280.66 pg/ml). Mean IL-10 prior to exposure was 286.35 pg/ml , which was not signifi-

cantly different ($T = -3.0$, $p = 0.25$, power = 20%) from mean IL-10 at time of sacrifice (236.07 pg/ml).

Viremia was available from only 6 of the 8 animals which were exposed to VEEV. Virus titers were highest in animals sacrificed at 48 hours than in those other animals of known viremia. Virus titers for the six animals were as follows: 7800 PFU per ml of blood for the two animals sacrificed 48 hours post-exposure; 370 PFU per ml for the two animals sacrificed 72 hours post-exposure; and 50 and 25 PFU per ml for the two animals sacrificed 96 hours post-exposure.

Pre-exposure testosterone and cortisol levels were both positively associated with viremia levels after exposure ($r = 0.98$, $p = 0.02$ and $r = 0.96$, $p = 0.04$, respectively; controlling for the hours between exposure and sacrifice; testosterone correlation controlling for cortisol levels, and vice versa).

The results of this preliminary study suggest that: 1) macaques infected with VEEV exhibit a febrile response with possibly few significant changes in cytokine levels; 2) higher levels of testosterone (when controlling for cortisol) prior to virus exposure may be directly associated with higher viremia after exposure; and 3) testosterone levels are lower during infection with VEEV than prior to infection.

First, macaques infected with VEEV exhibited a significant febrile response with few significant changes in cytokine levels. That is, temperature levels were elevated, IL-12 levels were lower, and IL-10 levels were relatively unchanged during infection. In both humans and non-human primates, VEEV infection has been characterized by fever with viremia and lymphopenia [7]. Unfortunately, no definitive conclusion can be drawn regarding the cytokine response to VEEV in the present study utilizing macaques because of the small sample size, lack of statistical power, and inability to assess the acute phase of infection (the sample was lacking information from animals at the 24 hour post-exposure timepoint). Furthermore, exposure to VEEV was not followed-up for longer than 96 hours, and

only a single sample was obtained from each animal throughout this short infection period.

Second, higher levels of testosterone prior to virus exposure were directly associated with higher viremia after exposure. That is, when controlling for time between exposure and sacrifice as well as cortisol levels, pre-exposure testosterone levels were significantly (directly) associated with viremia in post-exposure samples. In general, the effects of testosterone on viral infections, including HIV and Sindbis virus, are equivocal [11,14,15]. It may be the case that testosterone plays an immunosuppressive role in response to viral infection in male macaques and that higher levels of basal testosterone levels may increase an animal's susceptibility to viral infection. Future studies are warranted.

Thirdly, testosterone levels were significantly lower during infection with VEEV than prior to infection. Similarly, serum testosterone decreases during the onset of various conditions, including *Plasmodium vivax* infection in Honduran men [16], surgery and trauma in men [17], and *Trypanosoma brucei brucei* infection in rats [18]. Furthermore, testicular atrophy and azoospermia have been reported from men who died of AIDS [19], and azoospermia has been associated with SIV infection in young male rhesus macaques [20]. Depressed androgen levels during physiological perturbations may be an advantageous, adaptive host response in order to prevent immunosuppression by high testosterone levels and to re-direct energetic resources [21-23], specifically away from metabolically-expensive anabolic functions [11]. Although the results of the present study are very preliminary, they do suggest that disease-induced androgen suppression is a reasonable area for future study.

Abbreviations

Venezuelan equine encephalitis virus (VEEV), United States Army Medical Research Institute of Infectious Diseases (USAMRIID), plaque forming units (PFU), human immunodeficiency virus (HIV), acquired immunodeficiency syndrome (AIDS), simian immunodeficiency virus (SIV).

Competing interests

The author(s) declare that they have no competing interests.

Authors' contributions

MPM conducted the hormone analyses and wrote the manuscript. FBC edited the paper and coordinated the research efforts at Tulane Primate Center. MAJ edited the paper and conducted the cytokine analyses. JK conducted the animal work. GVL edited the paper and coordinated

the research efforts at USAMRIID. All co-authors read and approved the final manuscript.

Acknowledgements

The following individuals provided valuable logistical support: Richard Briescas, Susan Durham, Leann Myers, Michael Robich, Stephen Stearns, Brian Walker, and David Watts. Funding was provided by the Department of Bacteriology and Parasitology at Tulane National Primate Research Center, the Department of Tropical Medicine and Center for Infectious Diseases at Tulane University, and the Williams Fund and Graduate School Summer Study Fund from Yale University. Opinions expressed here do not necessarily reflect those of the Department of Defense.

References

1. Johnston RE, Peters CJ: **Alphaviruses**. In *Fields Virology* 3rd edition. Edited by: Fields BN, Knipe DM, Howley PM. New York: Raven Press; 1995:843-898.
2. Vogel P, Abplanalp D, Kell W, Ibrahim MS, Downs MB, Pratt WD, Davis KJ: **Venezuelan equine encephalitis in BALB/c mice: kinetic analysis of central nervous system infection following aerosol or subcutaneous inoculation**. *Arch Pathol Lab Med* 1996, **120**:164-721.
3. Charles PC, Trgovcich J, Davis NL, Johnston RE: **Immunopathogenesis and immune modulation of Venezuelan equine encephalitis virus-induced disease in the mouse**. *Virology* 2001, **284**:190-202.
4. Charles PC, Walters E, Margolis F, Johnston RE: **Mechanism of neuroinvasion of Venezuelan equine encephalitis virus in the mouse**. *Virology* 1995, **208**:662-671.
5. Bennett AM, Elvin SJ, Wright AJ, Jones SM, Philippotts RJ: **An immunological profile of BALB/c mice protected from airborne challenge following vaccination with a live attenuated Venezuelan equine encephalitis virus vaccine**. *Vaccine* 2002, **19**:337-347.
6. Grieder FB, Davis BK, Zhou XD, Chen SJ, Finkelman FD, Gause WC: **Kinetics of cytokine expression and regulation of host protection following infection with molecularly cloned Venezuelan equine encephalitis virus**. *Virology* 1997, **233**:302-312.
7. Pratt WD, Gibbs P, Pitt ML, Schmaljohn AL: **Use of telemetry to assess vaccine-induced protection against parenteral and aerosol infections of Venezuelan equine encephalitis virus in non-human primates**. *Vaccine* 1998, **16**:1056-1064.
8. Grossman CJ: **The role of sex steroids in immune system regulation**. In *Bilateral Communication between the Endocrine and Immune Systems* Edited by: Grossman CJ. New York: Springer-Verlag; 1995:1-11.
9. Chao TC, Van Alten PJ, Walter RJ: **Steroid sex hormones and macrophage function: modulation of reactive oxygen intermediates and nitrite release**. *Am J Reprod Immunol* 1994, **32**:43-52.
10. Daynes RA, Araneo BA: **Regulation of T-cell function by steroid hormones**. In *Cellular and Cytokine Networks in Tissue Immunity* Edited by: Meltzer MS, Mantovani A. New York: Wiley-Liss; 1991:77-82.
11. Muehlenbein MP, Bribiescas RG: **Testosterone-mediated immune functions and male life histories**. *Am J Hum Biol* 2005, **17**:527-558.
12. Elenkov IJ, Chrousos GP: **Stress, cytokine patterns and susceptibility to disease**. *Baillieres Best Pract Res Clin Endocrinol Metab* 1999, **13**:583-595.
13. Bambino TH, Hsueh AJ: **Direct inhibitory effect of glucocorticoids upon testicular luteinizing hormone receptor and steroidogenesis in vivo and in vitro**. *Endocrinology* 1981, **108**:2142-2148.
14. Lindstrom KM, Krakower D, Lundstrom JO, Silverin B: **The effects of testosterone on a viral infection in greenfinches (*Carduelis chloris*): an experimental test of the immunocompetence-handicap hypothesis**. *Proc R Soc Lond B Biol Sci* 2001, **268**:207-211.
15. Bhasin S, Storer TW, Javanbakht M, Berman N, Yarasheski KE, Phillips J, Dike M, Sinha-Hikim I, Shen R, Hays RD, Beall G: **Testosterone replacement and resistance exercise in HIV-infected men**

- with weight loss and low testosterone levels. *JAMA* 2000, **283**:763-770.
16. Muehlenbein M, Alger J, Cogswell F, James M, Krogstad D: **The reproductive endocrine response to *Plasmodium vivax* infection in Hondurans.** *Am J Trop Med Hyg* 2005, **73**:178-187.
 17. Spratt DI: **Altered gonadal steroidogenesis in critical illness: is treatment with anabolic steroids indicated?** *Baillieres Best Pract Res Clin Endocrinol Metab* 2001, **15**:479-494.
 18. Soudan B, Tetaert D, Racadot A, Degand P, Boersma A: **Decrease of testosterone level during an experimental African trypanosomiasis: involvement of a testicular LH receptor desensitization.** *Acta Endocrinol (Copenh)* 1992, **127**:86-92.
 19. Dym M, Orenstein J: **Structure of the male reproductive tract in AIDS patients.** In *Heterosexual Transmission of AIDS* Edited by: Alexander NJ, Gabelnick HL, Spieler JM. New York: Alan R. Liss; 1990:181-196.
 20. Nadler RD, Manocha AD, McClure HM: **Spermatogenesis and hormone levels in rhesus macaques inoculated with simian immunodeficiency virus.** *J Med Primatol* 1993, **22**:325-329.
 21. Folstad I, Karter AJ: **Parasites, bright males and the immunocompetence handicap.** *Am Nat* 1992, **139**:603-622.
 22. Sheldon BC, Verhulst S: **Ecological immunology: costly parasite defenses and trade offs in evolutionary ecology.** *TREE* 1996, **11**:317-321.
 23. Wedekind C, Folstad I: **Adaptive or nonadaptive immunosuppression by sex-hormones.** *Am Nat* 1994, **143**:936-938.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

